RLP
APPROVED

DISPLAYED BY INFOPEST

Product Name: DITRAC ALL WEATHER BLOX RODENTICIDE
APVMA Approval No: 49867/125129

Label Name:	DITRAC ALL WEATHER BLOX RODENTICIDE
Signal Headings:	POISON KEEP OUT OF REACH OF CHILDREN READ SAFETY DIRECTIONS BEFORE OPENING OR USING
Constituent Statements:	0.05 g/kg BRODIFACOUM contains Denatonium Benzoate (Bittering Agent)
Mode of Action:	
Statement of Claims:	Ready to use block bait for use in damp or dry situations in and around industrial, commercial, agricultural and domestic buildings. Controls species resistant to warfarin.
Net Contents:	1.8 - 10 kg Contains 20 g rodent blocks which are illegal to sell separately
Restraints:	DO NOT exceed 3 m between bait stations for mice, or 9 m for rats. DO NOT use bait stations further than 2 m from buildings. DO NOT apply baits to crops.
Directions for Use:	This section contains file attachment.
Other Limitations:	THE USE OF THIS PRODUCT FOR THE CONTROL OF PROTECTED NATIVE RODENTS REQUIRES PERMISSION FROM WILDLIFE AUTHORITIES.
Withholding Periods:	
Trade Advice:	

General Instructions:	GENERAL INSTRUCTIONS Locate bait stations along the walls or rafters and in dark sheltered spots in which there are signs of rodent activity. Nail or fix in place if relevant. Wax blocks are particularly suitable for use in damp situations where risk of contamination should be minimised. Record the number and locations of bait stations for the purpose of replenishment and later removal. Most deaths occur 4 to 7 days after consumption of bait.

Resistance Warning:	

Precautions:	PRECAUTION Keep away from food and water and surfaces which come in contact food and water.

Protections:	PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT DO NOT contaminate dams, streams, rivers, watercourses or drains with the bait or used containers. The bait is hazardous to pigs, cats, dogs, poultry, birds and wildlife and care should be exercised in its use. DO NOT place baits in locations which are accessible to domestic animals, livestock, nontarget native animals or birds. The product is restricted to use in and around buildings, (within 2 metres) or enclosed spaces.

Storage and Disposal:	Store in the closed, original container in a well-ventilated area, as cool as possible. DO NOT store for prolonged periods in direct sunlight. Store in a locked room or place away from children, animals, food and feedstuffs, seed and fertilisers. Do not dispose of chemicals on site. Break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Do not burn empty containers or product. Search for and burn or bury all rodent bodies, do not place in refuse bins or in rubbish tips. Ensure uneaten baits are removed and disposed of with the empty containers when the rodent activity has ceased.

Safety Directions:	Poisonous if swallowed. Do not inhale dust. Do not touch bait. Use scoop or measure. If on skin and after each baiting, wash thoroughly with soap and water. Vitamin K1 [phytomenadione] is antidotal.

First Aid Instructions:	If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26; New Zealand 0800 764 766.

First Aid Warnings:	

DIRECTIONS FOR USE

Baiting Strategy:

Eliminate as far as practicable, all alternative food sources. Bait areas where conditions might encourage rodent activity. Place baits under cover. Inspect bait stations regularly [during routine inspections]. Replace eaten or old/deteriorated baits. If quickly eaten increase the number of baits in the bait station. Continue observation and replenish until no more baits are taken. Baiting for at least two weeks is necessary to reduce rat/mouse numbers.

For mice: use single blocks. In very high infestation or fast uptake of the bait use 2 blocks.

For rats: use 1 to 3 blocks depending on level of infestation. In very high infestation or fast uptake use more baits per bait station.

SITUATION	PEST	RATE	CRITICAL COMMENTS
Agricultural, Commercial, Domestic, Industrial.	Rats (*Rattus norvegius, Rattus rattus*) Mice (*Mus musculus*)	1 to 3 blocks per bait station depending on level of infestation.	Set out baits in places frequented by mice and rats but not more than 2 m away from buildings etc. For rats: place 1 or more blocks per bait station at intervals of 5-9 m depending on level of uptake and infestation. For mice: place 1-2 blocks at 2-3 m intervals. In very high infestations and when using larger bait stations use more blocks per station and use the shorter distance between bait stations.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.